



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶:

C07K 5/037, A61K 38/04

A2

(11) International Publication Number:

WO 96/40739

(43) International Publication Date:

19 December 1996 (19.12.96)

(21) International Application Number: PCT/US96/09831

(22) International Filing Date: 7 June 1996 (07.06.96)

(30) Priority Data:

08/476,119

7 June 1995 (07.06.95)

US

(71) Applicant: TERRAPIN TECHNOLOGIES, INC. [US/US];
750-H Gateway Boulevard, South San Francisco, CA 94080 (US).

(72) Inventors: KAUVAR, Lawrence, M.; 1205 Cole Street, San Francisco, CA 94117 (US). LYTTLE, Matthew, H.; P.O. Box 1116, Point Reyes Station, CA 94956 (US). SATYAM, Apparao; 3545 Cade Drive, Fremont, CA 94536 (US).



(74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Foerster L.L.P., 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).

(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: URETHANE MEDIATED, GST SPECIFIC MOLECULAR RELEASE SYSTEMS



(57) Abstract

Compounds of formula (1) or of formula (2) or the amides, esters or salts thereof, wherein: S^x is $\text{S}=\text{O}$, $\text{O}=\text{S}=\text{O}$, $\text{S}=\text{NH}$, $\text{HN}=\text{S}=\text{O}$, $\text{Se}=\text{O}$, $\text{O}=\text{Se}=\text{O}$, $\text{Se}=\text{NH}$, $\text{HN}=\text{Se}=\text{O}$, S^+R^3 wherein R^3 is alkyl (1-6C) or $\text{O}-\text{C}=\text{O}$ or $\text{HN}-\text{C}=\text{O}$; each R^1 and R^2 is independently H or a noninterfering substituent; wherein (conj) represents a conjugated system capable of transmitting electrons; n is 0 or 1; YCO is selected from the group consisting of K-Glu, K-Glu-Gly, Glu, Glu-Gly, J-Asp, J-Asp-Gly, Asp and Asp-Gly; AA_C is an amino acid linked through a peptide bond to the remainder of said compound of formula (1); and $\text{N}(\text{Z})$ represents a reduced nitrogen-containing leaving group and L represents an electron-withdrawing leaving group, are useful as prodrugs and to generate active components released by the activity of glutathione S-transferase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

URETHANE MEDIATED, GST SPECIFIC MOLECULAR RELEASE SYSTEMS

Technical Field

The invention relates to compounds capable of releasing useful entities wherein the release is catalyzed by glutathione S-transferase (GST). More specifically, the invention concerns such compounds wherein the release is mediated by electron donation to the leaving group through a urethane linkage.

Background Art

Electron transit through a urethane linkage has been utilized to construct prodrugs by Senter, P.D. *et al.*, J Org Chem (1990), 55:2975. In this work, the reduction of a disulfide bridging two phenyl moieties was used to mediate the release of either nitroaniline or mitomycin C, wherein the amino group of the substance released was part of a urethane linkage para to the disulfide on one of the phenyl moieties. The reduction released electrons through the phenyl moiety to decompose the urethane. This provided the released nitroaniline or mitomycin C, and CO₂ as a by-product.

In addition, Nicolaou, K.C. *et al.*, Angew Chem Int Ed Engl (1991) 30:1032 describe the release of a dynemycin A analog wherein the amino group of the dynemycin A analog was included as part of a urethane linkage to the moiety f-SO₂CH₂OC(O)-N. Neither of these prodrug type molecules is enzyme regulated.

PCT/US1994/11109 referenced and incorporated above was published 13 April 1995 as WO95/09865. This published application describes a set of glutathione S-transferase activated compounds wherein the release of a desired leaving group is actuated by abstraction of a hydrogen ion H to the cysteinyl sulfur atom in a glutathione analog. The nature of the glutathione analog will determine which isoenzyme of GST will be the most effective in activating the release of the leaving group. Also disclosed in WO95/09865 is the inclusion of a urethane linkage within the leaving group, so that CO₂ is released when the leaving group is released as well.

The use of a conjugated X system to participate in the transfer of electrons from a relevant portion of a prodrug to the group released is also described by Papanastassiou,


Z. B. *et al.*, Experientia (1968) 24:325 and Tercel, M. *et al.*, J Med Chem (1993) 36:2578, as well as in the PCT reference described above.

It has now been found that the enzyme specificity conferred by the nature of glutathione analogs can be coupled with the electron release mechanisms associated with the urethane linkage to provide a new class of effective prodrugs for a variety of nitrogen-containing pharmaceuticals as well as more generally a release mechanism for any moiety containing reduced nitrogen. In addition, by taking advantage of the ability to move electrons through a conjugated system, the urethane-mediated linkage can be employed to release moieties which do not contain reduced nitrogen as part of the urethane linkage *per se*.

Disclosure of the Invention

The invention relates to a new class of GST-activated compounds capable of releasing desired moieties by electron transfer through a urethane linkage, with concomitant release of CO₂, optionally through coupling to a conjugated system. These compounds have the advantage of GST-regulated specificity, the equilibrium-driving properties associated with CO₂ release, and general applicability to release of electron withdrawing leaving groups. The compounds of the invention are therefore useful as prodrugs as well as laboratory reagents.

Thus, in one aspect, the invention is directed to a compound of the formula



or the amides, esters or salts thereof, wherein:

S^x is S=O, O=S=O, S=NH, HN=S=O, Se=O, O=Se=O, Se=NH, HN=Se=O,

S⁺R³ wherein R³ is alkyl (1-6C) or O-C=O or HN-C=O;

each R of R¹, and R² is independently H or a noninterfering substituent;

wherein (conj) represents a conjugated system capable of transmitting

electrons;


n is 0 or 1;

YCO is selected from the group consisting of K-Glu, K-Glu-Gly, Glu, Glu-Gly, JAsp, J-Asp-Gly, Asp and Asp-Gly;

AA_C is an amino acid linked through a peptide bond to the remainder of said compound of Formula 1; and

N(Z) represents a reduced nitrogen-containing leaving group.

The invention also relates to a compound of the formula



wherein S^x, R¹, R², YCO, conj, n, and AA_C are defined as above for Formula 1;

and

L represents an electron withdrawing leaving group.

In other aspects, the invention is directed to methods of synthesizing the compounds of Formulas 1 and 2, to pharmaceutical compositions containing these compounds, and to methods to impair or otherwise affect tumor cells or other targets by administering the compounds of Formulas 1 or 2 in contexts where the prodrugs are selectively cleaved by the targets to release N(Z) or L, which is typically a cytotoxic agent.

In still other aspects, the invention is directed to methods selectively to treat tumor cells or other target cells with characterized GST contents by selectively administering the prodrugs of the invention that are sensitive to cleavage with a GST that shows an elevated level in the target cells.


Modes of Carrying Out the Invention


The compounds of the invention are prodrugs which can be used selectively to target tissues having GST complements which are elevated or which contain isoenzymes peculiar in specificity to the prodrug provided. Depending on the nature of YCO and AA_C, these compounds are differentially activated by GST enzymes of the T, X and I classes. These prodrugs, in addition to being selective for cells with elevated GST complements *per se*, can be used in a finely tuned protocol to target cells which have elevated levels of a particular isoenzyme of the GST group.

In an additional use, the compounds of Formulas 1 and 2 can be used as analytical reagents for GST activity by employing as "L" or "N(Z)" an indicator group which is detectable when liberated from the compounds of Formulas 1 or 2. Such a reagent is suitable for determining the concentration of GST of known substrate specificity, or analyzing the specificity of particular GSTs by varying the glutathione analog component of the compounds of Formulas 1 or 2.

Compounds of the Invention

The compounds of the invention are comprised of a tripeptide which is glutathione or an analog thereof coupled to a leaving group through a molecular system which permits release of the leaving group N(Z) or L when the compounds of Formulas 1 or 2 are treated with the appropriate GST. CO₂ will also be released. The release of the leaving group occurs through a "J-elimination" -- i.e., the removal of the proton on the carbon I to the electron-poor oxidized carbon, sulfur or selenium releases electrons which are ultimately absorbed by the leaving group and result in its release. This can be shown schematically as follows:





The electron pair can be released to the leaving group through liberation of CO₂ directly through J-elimination as shown above or through a system of conjugation represented by (conj), when n is 1 in Formula 1 or 2.

The substituents R¹ and R² play no direct part in the release of substituent N(Z) or L and simply must be noninterfering substituents. The rate of J-elimination can be controlled by the nature of these R groups; by choosing electron withdrawing or electron donating substituents the rate of elimination can be accelerated or decreased. Suitable substituents for R¹ and R² include H, substituted or unsubstituted alkyl (1-6C) substituted or unsubstituted aryl (6-12C), substituted or unsubstituted aryl alkyl (7-12C), cyano, halo, substituted or unsubstituted alkoxy (1-6C), substituted or unsubstituted aryloxy (6-12C) or substituted or unsubstituted arylalkyloxy (7-12C).

Alkyl, aryl, and arylalkyl have their conventional meanings; alkyl groups are straight, branched chain or cyclic saturated hydrocarbon moieties such as methyl, tert-butyl, cyclohexyl, and the like. Aryl groups include aromatic systems such as phenyl, naphthyl, pyridyl and the like. Arylalkyl substituents contain an aryl moiety coupled to the remainder of the molecule through an alkylene moiety. Such groups include, most commonly benzyl, phenylethyl, 2-pyridylethyl, and the like.

Suitable substituents in the substituted forms include halo, SR, OR, and NR₂ wherein R is H or lower alkyl (1-4C).

Preferred embodiments for R¹ and R² independently are H, lower alkyl (1-4C) and phenyl. In particularly preferred embodiments, R¹ is H or phenyl, all R² are H and n=0. However, any noninterfering substituents may be used as R¹ and R². These substituents are independently embodied.

The embodiments of YCO and -AA_C determine the nature of the glutathione-like tripeptide. A preferred embodiment is that wherein YCO is K-glutamic and AA_C is

glycine, phenylglycine, J-alanine, alanine or phenylalanine, resulting in the tripeptide glutathione or a close analog. However, alternative embodiments of YCO include J-Asp, Glu, Asp, K-GluGly, J-AspGly, GluGly and AspGly. Alternative embodiments of AA_C include, along with the preferred glycine, phenylglycine, J-alanine, alanine, and unsubstituted phenylalanine: valine, 4-aminobutyric acid, aspartic, substituted phenylglycine, histidine, tryptophan, tyrosine, and substituted phenylalanine. Suitable phenylalanine and phenylglycine substituents are as described above for the substituted forms of R¹ and R².

Suitable embodiments for L include those which generate drugs which may be cytotoxic to unwanted cells. Such drugs include the phosphoramidate mustards, the phosphorodiamidate mustards, the chemotherapeutic agents adriamycin and daunorubicin, toxins such as ricin toxin or diphtheria toxin, antiinflammatory or steroid-based drugs and the like, and other metabolic modulators such as 2,3-di-t-butyl-4-hydroxyanisole. Preferred forms of the phosphorodiamidate mustards are -OP(O)(N(CH₂CH₂Cl)₂)₂, -OP(O)(N(CH₂CH₂Br)₂)₂, -OP(O)(NHCH₂CH₂Cl)₂ and -OP(O)(NHCH₂CH₂Br)₂. Any biologically active moiety, provided with an electron adsorbing linkage to the remainder of the compound so that "L" released by J-elimination may be used.

For embodiments of compounds of Formulas 1, the released moiety N(Z) includes, by definition, reduced nitrogen. Suitable released compounds then include the nitrogen mustards such as bis(2-chloroethyl)amine; uracil mustards wherein the bis (2-chloroethyl)amine is a substituent at the 5 position of the uracil ring and other mustards which include primary or secondary amines; various antibiotics which contain suitable amino groups to participate in the urethane linkage such as mitomycin C, actinomycin D, and the vinca alkaloids vincristine and vinblastine, and a dynemycin analog which subsequently follows a reaction path leading to a benzenoid diradical capable of effecting DNA inactivation by hydrogen atom abstraction.

As stated above, electron release can also be mediated through conjugated systems either to avoid the necessity for the inclusion of reduced nitrogen in the released moiety or simply to provide for electron flow or both. These conjugated systems may either be alkylene-based straight chain moieties such as -CR=CR-, -CR=CR-CR=CR-, -CR=CR-CR=CR-CR=CR- and the like, or may be included in aliphatic or aromatic ring systems such as 1,3,cyclohexadiene wherein the ring system is included in the compound

through bonds at the 1 and 4 positions, or benzene or other aromatic systems which are included through bonds to even numbers of carbons.

As shown in Formula 2, advantage may be taken of conjugated systems to liberate any moiety which can absorb electrons. For example, in an article by Mulcahy, R.T. *et al.*, J Med Chem (1994) 37:1610, liberation of a phosphoramidate mustard was described. The phosphoramidate is coupled through a methylene linkage to a para-nitrobenzene moiety and the conjugate is reduced under hypoxic conditions present in some cells to liberate the phosphoramidate mustard $OP(O)(N(CH_2CH_2Cl)_2)_2$ leaving behind a para-phenylene monoamine. In the present invention compounds, similar liberation of the phosphoramidate mustard through the mediation of the para-nitrogen is effected by electron donation through the urethane moiety to the aromatic ring, again resulting in the para-phenylene monoamine and phosphoramidate mustard.

Similarly, in a manner analogous to the known generation of the extremely cytotoxic nitrogen mustard mechlorethamine ($Me-N(CH_2CH_2Cl)_2$) by reduction of the quaternary amine coupled through a methylene linkage to ortho or para nitrobenzene, the J-elimination can again be used as a source of electrons through the mediation of a urethane linkage to generate the mechlorethamine and phenylene monoamine byproduct. Non-GST mediated hypoxic release from ortho or para nitrobenzyl is described by Papanastassiou, Z.B. *et al.*, Experientia (1968) 24:325; Tercel, M. *et al.*, J Med Chem (1993) 36:2578.

Preferred compounds of the invention are:

K-Glutamyl-I-amino-J(2-ethyl, N,N-bis(2'-chloroethyl)carbamoyl)sulfonyl)propionyl glycine;

K-Glutamyl-I-amino-J(2-ethyl, N,N-bis(2'-chloroethyl)carbamoyl)sulfonyl)propionyl phenyl glycine;

K-Glutamyl-I-amino-J-((2-ethyl-(4-benzyloxy(N,N,N',N' tetrakis(2-chloroethyl)phosphorodiamidate)) carbamido)sulfonyl)propionyl glycine;

K-Glutamyl-I-amino-J-((2-ethyl-(4-benzyloxy(N,N,N',N' tetrakis(2-chloroethyl)phosphorodiamidate)) carbamido)sulfonyl)propionyl phenyl glycine; and their diethyl esters.

In addition, indicator molecules such as p-nitrophenol can be used as leaving groups when the compounds of Formulas 1 or 2 is intended as a reagent.

The compounds of the invention may also be prepared in the forms of their esters or amides, or as their salts. The esters, amides or salts are formed with any or all carboxyl groups present in the molecule; hence, included in this group are monoesters, diesters, and, if applicable, triesters. Similarly, monoamides, diamides, or, if applicable, triamides are included.

The esters or amides may be alkyl (1-6C), alkenyl (1-6C) or arylalkyl (7-12C). Alkyl esters of the free carboxyls are esters of the straight- and branched-chain alkyl alcohols (1-6C) such as methanol, ethanol, isopropanol, t-butanol, n-hexanol and the like. Suitable alkyl (1-6C) amides are those of primary straight- or branched-chain alkyl amines, such as methylamine, ethylamine, n-propylamine, isopentylamine, and isohexylamine. Alkenyl esters are similar, but contain at least one double bond. Arylalkyl is as defined above. The alcohols or amines may also carry noninterfering substituents such as halo, alkoxy, or alkyl amines. The esters and amides are prepared using conventional techniques, with suitable protection of any alcohol or amino functional groups in the compound of Formula 1.

The salts of the compounds of the invention may be formed of inorganic or organic bases to form the basic salts of the free carboxyl groups or may be formed from organic or inorganic acids to obtain the acid addition salts of free amino groups. Thus, the salts may be of inorganic bases such as sodium hydroxide, potassium hydroxide, calcium hydroxide, ammonium hydroxide, magnesium hydroxide, and the like, or of organic bases such as trimethylamine, pyridine, pyrimidine, piperidine, lysine, caffeine, and the like. The acid addition salts may be formed from inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, and the like, or from organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, and the like. Salts of citric acid are preferred.

The salts of the compounds of Formula 1 are formed in standard protocols by treating with the appropriate base or acid at a temperature of from about 00C to about 1000C, preferably at room temperature either in water alone or in combination with an inert water-miscible organic solvent such as methanol, ethanol or dioxane.

Use of the Invention Compounds for Targeted Drug Delivery

The invention provides a general vehicle for delivering drugs to tissues specifically based on their GST content. The leaving group, when released in the target tissue, will exert its desired effects selectively in that target tissue. In addition to cytotoxicity, the released moiety may have other regulatory features. For example, where "L" is 2,3-di-t-butyl-4-hydroxyanisole, this compound is known to induce the synthesis of GSTs in mice. Administration of the compound of Formula 2 wherein "L" is 2,3-di-t-butyl-4-hydroxyanisole, will release this moiety may result in concomitant increase in GSTs. The target cells where release will occur can be regulated by manipulating the nature of the glutathione analog portion of the molecule. It may be desirable to enhance the GST component of the tumor cells concomitantly with supplying a compounds of Formulas 1 or 2 containing a cytotoxin.

As described above and demonstrated in the examples below, the various prodrugs of the invention are selective for the various isozymes of GST whose levels may be elevated in tumor cells. By determining the profile of GST isoenzyme levels in the tumor target, and matching this with the specificity of the prodrug, maximum effectiveness against the tumor cell will be obtained and maximum selectivity for the tumor cell as opposed to normal tissue can be achieved.

The compounds of Formulas 1 or 2 are administered as pharmaceutical compositions in usual formulations such as those outlined in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, latest edition. Typical formulations will include those for injection, for transdermal and transmucosal administration, and for oral administration. The formulations, depending on the intended mode, may be liquids, syrups, powders, capsules, suppositories, and the like. The compounds of the invention may be included in liposomes, or in other emulsified forms. Protocols for administration and suitable formulations are subject to optimization using standard procedures known to those in the art.

The antitumor activity of the invention compounds coupled with phosphorodiamidate mustard or other toxins can be assessed using a number of human tumor xenographs to determine tumor growth inhibition or a B16 mouse melanoma and measuring the prolongation of survival to determine the efficacy of particular compounds.

Compounds for Assay of GST Isoenzyme Activity

An alternative use for the compounds of Formulas 1 or 2 is as reagents in assays where the moiety "N(Z)" "L", when released from the compound can be readily detected. The compounds can thus conveniently be used to monitor the extent of the GST cleavage reaction, e.g., colorimetrically. Thus, an indicator moiety, such as p-nitrophenol, which is colorless when coupled to GSH or a GSH analog but develops a color on release from the compound by GST, offers an improved method of assaying GST activity. GST isoenzyme-specific assays using compounds comprising certain GSH analogs that are substrates only for selected GST isoenzymes can be used to determine substrate specificity.

Synthesis of the Invention Compounds

The compounds comprising glutathione or its analogs described above coupled to a desirable leaving group can be synthesized using means generally known in the art. Where S^x is an oxidized form of S or Se, the methods illustrated below can be used, incorporating modifications which render them applicable to desired compounds of the invention.


Thus, for example, compounds of Formulas 1 or 2 wherein S^x is $S=O$, $Se=O$, $O=S=O$ or $O=Se=O$ can be produced from the corresponding compounds wherein S^1 is S or Se, respectively, by oxidation with mild oxidizing agents such as peroxide or peracetate. Compounds of Formulas 1 or 2 wherein S^x is $S=NH$, $Se=NH$, $O=S=NH$, or $O=Se=NH$ can be obtained by treatment of the appropriate precursor, or a partially oxidized form, with chloramine T under conditions known in the art. Alternatively, the method of Whitehead, J.K. *et al.*, J Chem Soc (1952) 1572-1574, may be used. Precursor compounds lacking Y-CO or AA_C can be converted to the compounds of Formulas 1 or 2 by coupling the Y-CO moiety through a peptide linkage or the AA_C amino acid using standard peptide coupling techniques. When S^x is S or Se in reduced form in these precursors, these compounds may, similarly, be converted to compounds containing S or Se in oxidized form. Compounds of Formula 1 or 2 wherein S^x is a sulfonium ion, i.e., is S^+ , may be synthesized by treating compounds with reduced -S- with alkyl halides under suitable conditions to alkylate the sulfide. R^3 is alkyl

(1-6C) as defined above. Preferred alkyl halides for reaction to form, ultimately, compounds of Formulas 1 or 2 in this embodiment are the iodides.

For compounds of Formula 1 or 2 wherein S^x is $O-C=O$ are obtained using as a dipeptide or tripeptide starting material analogs of glutathione wherein serine substitutes for the cysteine moiety. Where S^x is $NH-C=O$, the corresponding amidation reaction is effected with analogs wherein 2,3-diaminopropionic acid replaces cysteine.


Preferred methods of synthesis are illustrated below. Reaction Scheme 1 shows the synthesis of a compound of Formula 1; the compound used for illustration is the urethane mustard of the oxidized (sulfone) of K-Glu-Cys-Glu; however, analogous pathways may be used to synthesize generally the class of compounds of Formula 1.

In the illustrative scheme shown, treatment of 2-bromoethyl chloroformate with dichlorodiethylamine in the presence of triethylamine yields the urethane bromide. Reaction of glutathione with this compound at pH 9-10 gives the glutathione conjugate, which is oxidized with hydrogen peroxide and peracetic acid to yield the sulfone of Formula 1.



Reaction Scheme 1

Reaction Scheme 2 shows the synthesis of an illustrative compound of Formula 2. As shown in Reaction Scheme 2, 2-bromoethyl chloroformate is reacted with phydroxymethylvaniline in triethanolamine and methylene chloride to provide the urethane bromide. The urethane bromide is then treated with POCl_3 and bis(2-chloroethyl)amine to give the tetrachloroethylphosphorodiamidate which is then treated first with glutathione or the relevant glutathione analog and then oxidized with peracetic acid to give the compound of Formula 2 as shown.



Reaction Scheme 2

The following example is intended to illustrate but not to limit the invention.

Example 1A Synthesis of the Diethyl Ester of the UrethaneMustard Conjugate of Oxidized K-Glu-Cys-GlyA. 2-Bromoethoxycarbonyl[bis(2-chloroethyl)amine]

2-Bromoethyl chloroformate (5.6 mL, 50 mmol) was added to a stirred suspension of bis(2-chloroethyl)amine hydrochloride (9.8 g, 55 mmol) in 250 mL of dry dichloromethane at 0-50C under argon over 2 min followed by 28 mL (200 mmol) of triethylamine over 20 min. The mixture was stirred at 5-100C for 3 h and at room temperature for 18 h, then suction filtered. The filtrate was concentrated *in vacuo*.

The residue was dissolved in 200 mL ethyl acetate and suction filtered to remove the triethylamine hydrochloride. The filtrate was successively washed with 100 mL each of 2N HCl, 5% NaHCO₃, water, and brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo* to give 13 g of crude product as a colorless oil, which was further purified by flash column chromatography (31-X 3.7-cm of silica gel bed and eluted isocratically with dichloromethane) to give 12.5 g (85%) of the title compound as a colorless oil: Anal. (C₇H₁₂BrCl₂NO₂) C, H, N.

B. K-Glutamyl-I-amino-J-[[2-ethoxycarbonyl-bis(2-chloreth-yl)-amine]-thio]propionylglycine

Glutathione (6.14 g, 20 mmol) was dissolved in 100 mL water and the pH was adjusted to between 9-10 by adding 1N NaOH. To this stirred solution at room temperature was added a solution of the urethane bromide prepared in paragraph A (2.93 g, 10 mmol) in 100 mL of 1:1 ethanol/acetonitrile. The resulting clear colorless solution was stirred at room temperature under argon for 3 days. TLC of the mixture indicated completion of the reaction.

The mixture was acidified to pH 5-6 with 10% acetic acid and most of the organic solvent portion was removed *in vacuo*. The aqueous portion was lyophilized and purified by HPLC: (buffer A, 0.1% TFA in 1:9 acetonitrile/water; buffer B, 0.1% TFA in 9:1 acetonitrile/water); eluted by running a gradient from 0-100% buffer B at an elution rate of 12 mL/min) to give 1.7 g (33%) of the title compound as a white fluffy hygroscopic powder: mp 85-1130C. Anal. (C₁₇H₂₈Cl₂N₄O₈S.TFA.2.5 H₂O) C, H, N.

C. K-Glutamyl-I-amino-J-[[2-ethoxycarbonyl-bis(2-chloroethyl)-amine]-sulfonyl]propionylglycine


To a stirred solution of the product of paragraph B (0.519 g, 1 mmol) in 10 mL glacial acetic acid at room temperature was added 30% H_2O_2 (0.39 mL, 2 mmol). The reaction flask was covered with aluminum foil to exclude light and the mixture was stirred at room temperature for 4 h. The mass spectrum indicated complete conversion to sulfoxide. 0.26 mL (1.25 mmol) of 32% peracetic acid in acetic acid was added to the mixture and it was stirred at room temperature for an additional 4 h, whereupon the mass spectral analysis of the mixture indicated formation of the title compound. The mixture was lyophilized and purified by HPLC to give 0.44 g (80%) of product as a hygroscopic white fluffy powder: mp 82-930C. Anal. ($\text{C}_{17}\text{H}_{28}\text{Cl}_2\text{N}_4\text{O}_{10}\text{S} \cdot \text{TFA} \cdot 2\text{H}_2\text{O}$) C, H, N.

D. K-Glutamyl-I-amino-J-[[2-ethoxycarbonyl-bis(2-chloroethyl)-amine]-sulfonyl]propionylglycine diethyldiester

To a stirred suspension of the product of paragraph C. (0.39 g, 0.7 mmol) in 28 mL of dry ethanol in a 100 mL reaction flask fitted with a reflux condenser under argon at room temperature was added thionyl chloride (1.1 mL, 15 mmol) from the top of the condenser. The resulting clear colorless solution was stirred at gentle reflux temperature for 2.5 h. The mass spectrum indicated formation of the diethyl diester. The mixture was concentrated *in vacuo* and the gummy residue was purified by HPLC to give 0.17 g (40%) of the title compound as a hygroscopic fluffy white powder: mp 54-600C. Anal. ($\text{C}_{21}\text{H}_{36}\text{Cl}_2\text{N}_4\text{O}_{10}\text{S} \cdot \text{HCl}$) C, H, N.

Claims

1. A compound of the formula



or the amides, esters or salts thereof, wherein:

S^x is $\text{S}=\text{O}$, $\text{O}=\text{S}=\text{O}$, $\text{S}=\text{NH}$, $\text{HN}=\text{S}=\text{O}$, $\text{Se}=\text{O}$, $\text{O}=\text{Se}=\text{O}$, $\text{Se}=\text{NH}$, $\text{HN}=\text{Se}=\text{O}$, S^+R^3 wherein R^3 is alkyl (1-6C) or $\text{O}-\text{C}=\text{O}$ or $\text{HN}-\text{C}=\text{O}$;

each R of R^1 , and R^2 is independently H or a noninterfering substituent;

wherein (conj) represents a conjugated system capable of transmitting electrons;

n is 0 or 1;

YCO is selected from the group consisting of K-Glu, K-Glu-Gly, Glu, Glu-Gly, J-Asp, J-Asp-Gly, Asp and Asp-Gly;

AA_C is an amino acid linked through a peptide bond to the remainder of said compound of Formula 1; and

$\text{N}(\text{Z})$ represents a reduced nitrogen-containing leaving group.

2. The compound of claim 1 wherein n is 0 and/or

wherein YCO is K-glu; and/or


wherein all of R^1 and R^2 are H; and/or

wherein S^x is $\text{O}=\text{S}=\text{O}$; and/or

wherein AA_C is selected from the group consisting of glycine, phenylglycine, J-alanine, alanine and phenylalanine; and/or

wherein $\text{N}(\text{Z})$ is selected from the group consisting of bis(2-chloroethyl)amine, a uracil mustard, mitomycin C, actinomycin C, vincristine, vinblastine, and dynemycin A.

3. A compound of the formula



or the amides, esters or salts thereof, wherein:

S^x is $\text{S}=\text{O}$, $\text{O}=\text{S}=\text{O}$, $\text{S}=\text{NH}$, $\text{HN}=\text{S}=\text{O}$, $\text{Se}=\text{O}$, $\text{O}=\text{Se}=\text{O}$, $\text{Se}=\text{NH}$, $\text{HN}=\text{Se}=\text{O}$,

S^+R^3 wherein R^3 is alkyl (1-6C) or $\text{O}-\text{C}=\text{O}$ or $\text{HN}-\text{C}=\text{O}$;

each R of R^1 , and R^2 is independently H or a noninterfering substituent;

wherein (conj) represents a conjugated system capable of transmitting electrons;

n is 0 or 1;

YCO is selected from the group consisting of K-Glu, K-Glu-Gly, Glu, Glu-Gly, JAsp, J-Asp-Gly, Asp and Asp-Gly;

AA_C is an amino acid linked through a peptide bond to the remainder of said compound of Formula 1; and

L represents an electron withdrawing leaving group.

4. The compound of claim 3 wherein n is 0 and/or

wherein YCO is K-glu; and/or

wherein all of R^1 and R^2 are H; and/or

wherein S^x is $\text{O}=\text{S}=\text{O}$; and/or

wherein AA_C is selected from the group consisting of glycine, phenylglycine, J-alanine, alanine and phenylalanine; and/or

wherein (conj) is para-phenylene; and/or

wherein L is a phosphoroamide mustard, a phosphorodiamidate mustard, adriamycin or daunorubicin.

5. A pharmaceutical composition for drug delivery which composition comprises as active ingredient the compound of claim 1 in admixture with a pharmaceutically acceptable excipient.

6. A pharmaceutical composition for drug delivery which composition comprises as active ingredient the compound of claim 3 in admixture with a pharmaceutically acceptable excipient.

7. A method to deliver a biologically active moiety to a target which method comprises administering to a subject containing said target the compound of claim 1 or a pharmaceutical composition thereof.

8. A method to deliver a biologically active moiety to a target which method comprises administering to a subject containing said target the compound of claim 3 or a pharmaceutical composition thereof.

PCT



WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/037, A61K 38/04	A3	(11) International Publication Number: WO 96/40739 (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09831 (22) International Filing Date: 7 June 1996 (07.06.96) (30) Priority Data: 08/476,119 7 June 1995 (07.06.95) US (71) Applicant: TERRAPIN TECHNOLOGIES, INC. [US/US]; 750-H Gateway Boulevard, South San Francisco, CA 94080 (US). (72) Inventors: KAUVAR, Lawrence, M.; 1205 Cole Street, San Francisco, CA 94117 (US). LYTTLE, Matthew, H.; P.O. Box 1116, Point Reyes Station, CA 94956 (US). SATYAM, Apparao; 3545 Cade Drive, Fremont, CA 94536 (US). (74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Foerster L.L.P., 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 30 January 1997 (30.01.97)	

(54) Title: URETHANE MEDIATED, GST SPECIFIC MOLECULAR RELEASE SYSTEMS



(57) Abstract

Compounds of formula (1) or of formula (2) or the amides, esters or salts thereof, wherein: S^x is S=O, O=S=O, S=NH, HN-S=O, Se=O, O=Se=O, Se=NH, HN=Se=O, S⁺R³ wherein R³ is alkyl (1-6C) or O-C=O or HN-C=O; each R of R¹, and R² is independently H or a noninterfering substituent; wherein (conj) represents a conjugated system capable of transmitting electrons; n is 0 or 1; YCO is selected from the group consisting of K-Glu, K-Glu-Gly, Glu, Glu-Gly, JAsp, J-Asp-Gly, Asp and Asp-Gly; AA_c is an amino acid linked through a peptide bond to the remainder of said compound of formula (1); and N(Z) represents a reduced nitrogen-containing leaving group and L represents an electron-withdrawing leaving group, are useful as prodrugs and to generate active components released by the activity of glutathione S-transferase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INTERNATIONAL SEARCH REPORT

Inte- Application No
PCT/US 96/09831

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K5/037 A61K38/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 09866 A (TERRAPIN TECH INC) 13 April 1995 see page 7, line 29 - page 8, line 18; claims; examples -----	1-8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

14 November 1996

Date of mailing of the international search report

06.12.96

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fuhr, C

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 96/09831

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 7 and 8
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/09831

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9509866	13-04-95	US-A- 5545621	13-08-96
		US-A- 5556942	17-09-96
		AU-A- 7962394	01-05-95
		AU-A- 8072094	01-05-95
		CA-A- 2173130	13-04-95
		EP-A- 0721465	17-07-96
		WO-A- 9509865	13-04-95
